

Characterization of New Bacteriophages of *Salmonella* and *Shigella* For Polivalent Intestinal “Enco Bacterophage” Preparation

Gabisonia T^{*}, Giorgadze I, Topuria-Nakaidze N, Loladze M, Chakhunashvili N, Nadiradze M, Eliava T, Kalandarishvili T, Tamarashvili N and Katamadze T

G. Eliava Institute of Bacteriophages, Microbiology and Virology, Tbilisi, Georgia.

*Correspondence:

Gabisonia T, Doctor of Science, Head, G. Eliava Institute of Bacteriophages, Microbiology and Virology, Tbilisi, Georgia, Tel: +995 577 423225.

Received: 16 March 2019; Accepted: 04 April 2019

Citation: Gabisonia T, Giorgadze I, Topuria-Nakaidze N, et al. Characterization of New Bacteriophages of *Salmonella* and *Shigella* For Polivalent Intestinal “Enco Bacterophage” Preparation. *Microbiol Infect Dis*. 2019; 3(2): 1-3.

ABSTRACT

New, highly active clones of bacteriophage against *S. typhimurium*, *S. sonnei* and *S. flexneri* strains have been isolated. Susceptibility of *S. typhimurium* strains to the phages made up 96.3%. The phage adsorption period on the host cell varies within 10-15 min, the latent period is 22-28 min, and the yield of the phage makes up 110-180 phage particles per virion. *Salmonella* phages belong to the morphological types of Myoviridae and Siphoviridae, while *shigella* phages belong to the morphological types of Siphoviridae, Podoviridae and Myoviridae.

Keywords

Salmonellae, *Shigella*, Bacteriophage, Morphology, Biological properties.

Introduction

Salmonella is important food-borne pathogen that causes food poisoning. Salmonellosis is a major illness accompanied by headache, diarrhea, vomiting, and high fever due to *Salmonella* infection in the epithelial tissue of animals and humans via contaminated food [1]. *Shigella* is another wide-spread food-borne pathogen causing human diarrhea termed “shigellosis” [2]. These bacteria are responsible for food poisoning outbreaks worldwide through various foods, such as ground meat, raw milk, apple cider, and fresh vegetables. In the United States, approximately 500,000 cases of shigellosis and more than 1.4 million cases of food-borne salmonellosis have been reported per year, with 17,000 hospitalizations and 600 deaths [3,4]. Wide and not always justified use of antibiotics and chemo preparations in medicine, especially of a broad spectrum, promoted the ubiquitous spread of bacteria with natural and acquired resistance. At the same time, intensive antibiotic therapy is often accompanied by a number of complications – changes in the immune reactivity, development of infections caused by opportunistic microorganisms.

Even though various antibacterial preparations have been

developed to control these pathogens, development of safe and effective new agents to control food-borne pathogens is urgently needed [5].

Existing in nature abundance of bacteriophage clones with different morpho-biological properties and molecular organization, differing from one another both in mechanism of interaction with host cells and reproductive capacity, requires a clear differentiation and in-depth study of principal and some subsidiary taxonomic characters for their purposeful use in the treatment and prevention of bacterial infection.

The aim of this work was to study the morphological and some biological properties of new phage clones (pure lines) of *salmonella* and *shigella* phages.

Materials and Methods

Isolation of bacteriophages from wastewater. 10 ml of concentrated broth is added to 90 ml of wastewater, and to isolate a particular phage, 1 ml of the corresponding 18-hour culture is added to this mixture [6].

After that, the whole mixture is placed in a thermostat at 37°C for 24 h. After 18-24 h of incubation, material is filtered through Millipore filters and checked for the presence of phage.

The study of a number of theoretical and practical problems requires availability of pure phage lines. For this purpose, the cloning method is used. Cloning of bacteriophages is carried out by passage of morphologically homogeneous negative colonies on homologous bacterial cultures. To study the interaction of the phage with the host cell, adsorption and phage yield are examined. To study the morphology of the phage itself, an electron microscope is used.

Results and Discussion

Isolation of new phages of salmonella was conducted from wastewater, the Kura River and the stool of patients with salmonellosis. Five phages, conditionally named as S-25, S-146, SK-18, SS-19, S-171, were isolated. 110 strains of *S. typhimurium* were isolated from different clinics in Tbilisi (Republican Children's Hospital, Republican Infectious Diseases Hospital, Republican Hospital).

The susceptibility of freshly isolated strains to new bacteriophages was determined. As the results showed, 96.3% of *S. typhimurium* strains were sensitive to 5 new phages. To characterize the biological properties of *S. typhimurium* bacteriophages, the following bacterial features were used in the study: the morphology of negative colonies, the spectrum of the lytic action, the rate of adsorption and the average yield per one cell.

On the lawn, *S. typhimurium* phage culture gave two types of negative colonies: medium – from 1.5 to 2.0 mm in diameter with smooth edges and a transparent center and large – from 1.5 to 4.0 mm in diameter with a transparent center and even edges.

The adsorption time of the studied phages varied within 11-14 min, the latent period was 22-28 min, and the yield was 120-130 phage particles per virion.

It has been established that *S. typhimurium* phages have a binary type of symmetry, a head and a tail. They belong to the morphological types of *Myoviridae* and *Siphoviridae*. The head of phages is characterized by hexagonal symmetry, the dimensions of the phage head are different (Table 1). The main differences concern the structure of the tail.

Name of the phage	Host strain	Morphological group	Dimensions of the phage head		Size of the phage tail	
			Length × width	Length × width	Length × width	Length × width
<i>S. typhimurium</i> S-25	<i>S. typhimurium</i> Q	<i>Myoviridae</i>	60 nm × 50 nm	95 nm × 20 nm		
<i>S. typhimurium</i> S-146	<i>S. typhimurium</i> Q	<i>Myoviridae</i>	60 nm × 50 nm	95 nm × 15 nm		
<i>S. typhimurium</i> SK-18	<i>S. typhimurium</i> Q	<i>Myoviridae</i>	75 nm × 70 nm	10 nm × 15 nm		
<i>S. typhimurium</i> SS-19	<i>S. typhimurium</i> Q	<i>Siphoviridae</i>	45 nm × 45 nm	125 nm × 15 nm		
<i>S. typhimurium</i> S-171	<i>S. typhimurium</i> Q	<i>Siphoviridae</i>	45 nm × 45 nm	100 nm × 15 nm		

Table 1: Electron microscopic parameters for 5 phages of *S. typhimurium*.




Figure 1: Plaques of phage *S. typh.* S-25 on the strain *S. typhimurium* Q plaque's diameter is 1-2 mm.




Figure 2: *S. typh.* SK-18 and *S. typh.* SS-19 phages electron micrograph.

The isolation of new phages against *S. sonnei* and *S. flexneri* strains was carried out from wastewater as well as from the Kura River. Six phages in all were isolated, 4 phage clones against *S. flexneri* and 2 clones against – *S. sonnei*.

Phage clones were characterized by the following parameters: morphology of negative colonies, adsorption rate, and average yield per one cell. On the host strain, *S. flexneri* phage clones gave negative colonies from 4 to 5 mm in size, and *S. sonnei* phage clones – from 4 to 6 mm, with a transparent center and even edges.

The adsorption time for the studied clones of *S. flexneri* phages varied from 10 to 15 min, the latent period was 22-26 min, and the yield was 110-180 phage particles per virion.

For *S. sonnei* phage clones, the adsorption time varied from 10 to 15 min, the latent period was 22-26 min, and the yield was 110-180 phage particles per virion.

It is established that the Shigella phages have a binary type of symmetry, a head and a tail. They belong to the morphological types of *Siphoviridae*, *Podoviridae* and *Myoviridae*.

The head of phages is characterized by hexagonal symmetry, the

dimensions of the phage head are different (Table 2). The main differences concern the structure of the tail.

Phage name	Host strain	Morphological group	Dimensions of the phage head	Size of the phage tail
			Length × width	Length × width
<i>S. flexneri</i> 8	<i>S. flexneri</i> 2a 25	<i>Siphoviridae</i>	50 nm × 50 nm	140 nm × 15 nm
<i>S. flexneri</i> 3a 25	<i>S. flexneri</i> 2a 25	<i>Siphoviridae</i>	45 nm × 45 nm	125 nm × 15 nm
<i>S. flexneri</i> 7	<i>S. flexneri</i> 2a 25	<i>Siphoviridae</i>	50 nm × 50 nm	140 nm × 15 nm
<i>S. flexneri</i> 3	<i>S. flexneri</i> 1B	<i>Podoviridae</i>	50 nm × 50 nm	15 nm × 10 nm
<i>S. sonnei</i> 442	<i>S. sonnei</i> 1148	<i>Myoviridae</i>	60 nm × 60 nm	95 nm × 15 nm
<i>S. sonnei</i> 3	<i>S. sonnei</i> 32	<i>Podoviridae</i>	45 nm × 45 nm	20 nm × 10 nm

Table 2: Electron microscopic parameters for the 6 phages of *S. sonnei* and *S. flexneri*.




Figure 3: *Sh. flexneri* phage 7 and *Shigella flexneri* phage 3 electron micrograph.

Based on the data obtained, it can be concluded that the isolated new *S. typhimurium* phages are highly active against the strains of *S. typhimurium* – 96.3%. The isolated *S. typhimurium* phages belong to the morphological groups of Myoviridae and Siphoviridae. The isolated *S. sonnei* and *S. flexneri* phages belong to the morphological groups of *Siphoviridae*, *Podoviridae* and *Myoviridae*. They have the same adsorption time of 10-15 min, the latent period of 22-26 minutes and the yield of 110-180 phage particles per virion.

References

1. Barbara G, Stanghellini V, Berti-Ceroni C, et al. Role of antibiotic therapy on long-term germ excretion in faeces and digestive symptoms after Salmonella infection. *Aliment Pharmacol Ther.* 2000; 14: 1127-1131.
2. Heyn Lee, Hye-Jin Ku, Dong-Hoon Lee, et al. Characterization and Genomic Study of the Novel Bacteriophage HY01 Infecting Both *Escherichia coli* O157:H7 and *Shigella flexneri*: Potential as a Biocontrol Agent in Food. *PLoS One.* 2016; 11: e0168985.
3. Scallan E, Hoekstra RM, Angulo FJ, et al. Foodborne illness acquired in the United States--major pathogens. *Emerg Infect Dis.* 2011; 17: 7-15.
4. Mead PS, Slutsker L, Dietz V, et al. Food-related illness and death in the United States. *Emerg Infect Dis.* 1999; 5: 607-625.
5. Minjung Park, Ju-Hoon Lee, Hakdong Shin, et al. Characterization and Comparative Genomic Analysis of a Novel Bacteriophage, SFP10, Simultaneously Inhibiting both *Salmonella enterica* and *Escherichia coli* O157:H7. *Appl Environ Microbiol.* 2012; 78: 58-69.
6. Adams M. Bacteriophages. 1954.